

Warped cones and Haagerup property

Guoqiang Li and Xianjin Wang

Abstract

In this paper we show that the fibred coarse embeddability of a warped cone implies the Haagerup property of the appropriate group. Moreover, Kazhdan's property (T) of the group implies geometric property (T) of the warped cone.

1 Warped cones

Warped cones were introduced by John Roe in [1], where serve as the examples of coarse spaces which do not have property A or which are not coarsely embeddable into Hilbert space.

Let Y be a smooth compact manifold, or a finite simplicial complex. The open cone $X = \mathcal{O}Y$ may be defined in the following way: Embed Y smoothly in a high-dimensional sphere S^{N-1} , and let X be the union of all the rays through the origin in $\mathbb{R}^N$ that meet the embedded copy of Y , equip X with the metric induced from $\mathbb{R}^N$. Observe that we can parametrise points on $\mathcal{O}Y$ as pairs (y, t) , where $y \in Y$, $t \in \mathbb{R}^+$. Let G be a group acting on X . In general the maps $x \mapsto gx$ will not be translations. The basic idea of warping is to modify the metric on X so that the action of G becomes an action by translations, i.e. the distance from x to gx is uniformly bounded in X for each $g \in G$.

Definition 1.1. (See [1]) Let (X, d) be a metric space and G be a group acting by homeomorphisms on X , provided with a finite generating set S . The warped metric d_G on X is the greatest metric satisfying the following inequalities:

$$d_G(x, x') \leq d(x, x'), \quad d_G(x, sx) \leq 1, \quad \forall s \in S.$$

It is easy to show that $d_G(x, y)$ exists for all x, y and defines a metric on X , so is well-defined. Clearly, d_G captures the aim of making G act by translations. Moreover, the distance from x to gx depends on the size of g in G , so the group structure does have some effect on the space (X, d_G) . We will need the following more concrete characterisation of d_G :

Proposition 1.2. (See [1]) Let $x, y \in X$. For $g \in G$, let $|g|$ denote the word length of g relative to the generating set S . The warped distance from x to y is the infimum of all sums

$$\sum_{i=1}^{N-1} d(g_i x_i, x_{i+1}) + |g_i|$$

taken over all finite sequences $x = x_0, x_1, \dots, x_N = y$ in X and $g_0, \dots, g_{N-1}$ in G . Moreover, if $d_G(x, y) \leq k$, then one can find a finite sequence as above with $N = K$ such that the infimum is attained.

Proof. The above does define a metric, say *delta*, and it satisfies the conditions mentioned above, whence $\delta(x, y) \leq d_G(x, y)$ for all $x, y \in X$. Conversely, $d_G \leq \delta$ by assumptions on d_G and the triangle inequality.

Looking now at the second part, note that if $d_G(x, y) \leq k$, then there must exist sequence of the above type where at most k g_i are non-identity elements; as removing any terms where $g_i = e$ can only decrease the size of the sum, we can assume that any such sequence has at most k terms. For fixed endpoints x, y the space of such chains can be assumed to be a closed subspace of $B_X(x, d(x, y) + 1)^k \times B_G(e, k)^k$, which is compact by properness of X, G . Here the infimum is attained. $\square$

Remark 1.3. The warped metric is a proper metric. The coarse structure induced by warped metric does not depend on the choice of generating set S for G , nor on the choice of metric d within the coarse structure of X .

We can now make the main definition of warped cone.

Definition 1.4. (See [1]) Let Y be a smooth compact manifold, or a finite simplicial complex, and let G be a finitely generated group acting by homeomorphisms on Y . The group acts on $\mathcal{O}Y$ by $g : (y, t) \rightarrow (gy, t)$. The warped cone $\mathcal{O}_G Y$ is the coarse space obtained by warping $\mathcal{O}Y$ equipped with the metric d_G arising from the action.

2 Characterization of the Haagerup property

The follow result is essentially proved in [1]. For the convenience of the reader, we include the proof below.

Theorem 2.1. (See [1]) Let Γ be a compact Lie group, and let $G \leq \Gamma$ be a dense finitely generated subgroup, which we consider as a discrete group acting on Γ by left translation. If the warped cone $\mathcal{O}_G \Gamma$ is coarsely embeddable in Hilbert space. Then G has the Haagerup property.

Proof. Let $X = \mathcal{O}_G\Gamma$. To say that X admits a coarse embedding into Hilbert space is to say that X admits a continuous negative type kernel k which is effective, that is, there are unbounded increasing functions $\rho_1, \rho_2 : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that

$$\rho_1(d(x, x')) \leq k(x, x') \leq \rho_2(d(x, x')).$$

Averaging over Γ , we may assume that k is Γ -invariant. For a fixed g ,

$$h_t(g) = k((y, t), (gy, t))$$

depends only on g and is bounded by $\rho_2(|g|)$; so we can find a sequence $t_k \rightarrow \infty$ such that the limit

$$h(g) = \lim_k h_{t_k}(g)$$

exists. It is a negative type function and satisfies $h(g) \geq \rho(|g|)$, so it is proper. Thus G admits a proper negative type function, which has the Haagerup property. $\square$

Remark 2.2. In particular, if G has property (T), the $\mathcal{O}_G\Gamma$ has geometry property (T). Indeed, if G has property (T), then $\mathcal{O}_G\Gamma$ cannot be coarsely embedded in Hilbert space. Therefore, $\mathcal{O}_G\Gamma$ cannot be amenable. Due to Corollary 6.1 in [2], we know that $\mathcal{O}_G\Gamma$ has geometric property (T).

In the original work, the warped cone was required to be coarsely embeddable into Hilbert space. It is worth noting that a similar proof shows that the same hypothesis, with ‘fibred coarsely embeddable into Hilbert space’ replacing ‘uniformly embeddable Hilbert space’, implies that G has the Haagerup property.

Theorem 2.3. *Let Y be a compact metric space, and G a finitely generated group which acts on Y by left translation. Assume that Y is equipped with a finite G -invariant Borel measure. If $\mathcal{O}_GY$ admits a fibred coarse embedding into Hilbert space, then G has Haagerup property.*

Proof. To say that $X = \mathcal{O}_GY$ is fibred coarse embedding into Hilbert space is to say that there exist a field of Hilbert spaces $(H_x)_{x \in X}$ over X , a section $s : X \rightarrow \bigsqcup_{x \in X} H_x$ (that is, $s(x) \in H_x$), and two non-decreasing functions ρ_1 and ρ_2 from $[0, \infty)$ to $(-\infty, \infty)$ with $\lim_{r \rightarrow \infty} \rho_i(r) = \infty$ ($i = 1, 2$) such that, for any $r > 0$, there exists a bounded subset $K_r \subset X$ for which there exists a trivialization

$$t_C : (H_x)_{x \in C} \rightarrow C \times H$$

for each subset $C \subset X \setminus K_r$ of diameter less than r , that is, a map from $(H_x)_{x \in C}$ to the constant field $C \times H$ over C such that the restriction of t_C to the fibre H_x ($x \in C$) is an affine isometry $t_C(x) : H_x \rightarrow H$, satisfying the following:

(1) for any $x, x' \in C$, $\rho_1(d_G(x, x')) \leq \|t_C(x)(s(x)) - t_C(x')(s(x'))\| \leq \rho_2(d_G(x, x'))$;
(2) for any two subset $C_1, C_2 \subset X \setminus K_r$ of diameter less than r with $C_1 \cap C_2 \neq \emptyset$, there exists an affine isometry $t_{C_1 C_2} : H \rightarrow H$ such that $t_{C_1}(x) \circ t_{C_2}^{-1}(x) = t_{C_1 C_2}$ for all $x \in C_1 \cap C_2$.

For a fixed $r > 0$, for any $x, x' \in X \setminus K_r$ with $d_G(x, x') < r$, choose a subset $C \subset X \setminus K_r$ of diameter less than r containing x and x' , and define

$$K_r(x, x') = \|t_C(x)(s(x)) - t_C(x')(s(x'))\|^2$$

If C_1 and C_2 are two subset of $X \setminus K_r$ of diameter less than r both of which contain x and x' , t_{C_1} and t_{C_2} are the corresponding trivializations, it follows from (2) above that $t_{C_1}(x) \circ t_{C_2}^{-1}(x) = t_{C_1}(x') \circ t_{C_2}^{-1}(x')$. So

$$\begin{aligned} & \|t_{C_1}(x)(s(x)) - t_{C_1}(x')(s(x'))\|^2 \\ &= \|t_{C_1}(x') \circ t_{C_2}^{-1}(x') \circ t_{C_2}(x)(s(x)) - t_{C_1}(x) \circ t_{C_2}^{-1}(x) \circ t_{C_2}(x')(s(x'))\|^2 \\ &= \|t_{C_1 C_2} \circ t_{C_2}(x)(s(x)) - t_{C_1 C_2} \circ t_{C_2}(x')(s(x'))\|^2 \\ &= \|t_{C_2}(x)(s(x)) - t_{C_2}(x')(s(x'))\|^2 \end{aligned}$$

That is, the definition of $K_r(x, x')$ for $x, x' \in X \setminus K_r$ with $d_G(x, x') < r$ does not depend on the choice of the subset C and the trivialization t_C . Define $K_r(x, x') = 0$ if $d_G(x, x') \geq r$. Then K_r is a well-defined, r -locally conditionally negative definite kernel on $X \setminus K_r \times X \setminus K_r$. It follows from (1) above that, for any $x, x' \in X \setminus K_r$ with $d_G(x, x') < r$,

$$(\rho_1(d_G(x, x')))^2 \leq K_r(x, x') \leq (\rho_2(d_G(x, x')))^2$$

Let μ be a (normalised) G -invariant measure on Y , parametrise points on $\mathcal{O}_G Y$ as pair (y, t) . For all sufficiently large t , the warped distance from (y, t) to (gy, t) is precisely the word length of g . Thus, we can define a function $\phi_{r,t}$ on the r -ball $B_r = \{g \in G : |g| < r\}$,

$$\phi_{r,t}(g) = \int_Y K_r((y, t), (gy, t)) d\mu(y)$$

and define $\phi_{r,t} \equiv 0$ outside the r -ball of G . Note that $\phi_{r,t}$ is an r -ball conditionally negative definite function on G . Indeed, for any $g_1, g_2, \dots, g_n$ in G with $d(g_i, g_j) < r$ and real scalars $\lambda_1, \lambda_2, \dots, \lambda_n$ with $\sum_{i=1}^n \lambda_i = 0$, one has

$$\begin{aligned} \sum_{i,j} \lambda_i \lambda_j \phi_{r,t}(g_i^{-1} g_j) &= \sum_{i,j} \lambda_i \lambda_j \int_Y K_r((y, t), (g_i^{-1} g_j y, t)) d\mu(y) \\ &= \int_Y \sum_{i,j} \lambda_i \lambda_j K_r((y, t), (g_i^{-1} g_j y, t)) d\mu(y) \\ &\leq 0. \end{aligned}$$

And

$$\mu(Y)(\rho_1(d_G(x, x')))^2 \leq \phi_{r,t}(g_1^{-1}g_2) \leq \mu(Y)(\rho_2(d_G(x, x')))^2$$

for $g_1, g_2 \in G$ with $d(g_1, g_2) < r$. Hence, for a fixed t , letting $r \rightarrow \infty$, there exist a subsequence of the sequence $\{\phi_{r,t}\}_{r>0}$ which converges pointwise to a function $\phi_t : G \rightarrow \mathbb{R}$ that is conditionally negative definite on the whole G and bounded by $\mu(Y)(\rho_2(|g|))^2$, so we can find a subsequence $t_m \rightarrow \infty$ such that the limit

$$\phi(g) = \lim_{m \rightarrow \infty} \phi_{t_m}(g)$$

exists. It is conditionally negative definite, and proper since we have

$$\mu(Y)(\rho_1(|g|))^2 \leq \phi(g) \leq \mu(Y)(\rho_2(|g|))^2$$

for all $g \in G$. Hence, G has the Haagerup property. $\square$

References

- [1] J. Roe, *Warped cones and property A*, Geom. Topol. **9**(2005),163-178.
- [2] R. Willett, G. Yu. *Geometric property (T)*, Chinese Annals of Mathematics, Series B, **35**(2014),761-800.